

December 2018 -
Newsletter

Inside this Newsletter:

- Inflammatory Skin Disorders
- Hormone Testing



Thank you for your support!

Our last edition of Pathology Focus for 2018

On behalf of Australian Clinical Labs, we would like to begin this end-of-year edition of Pathology Focus by wishing you a safe and happy holiday season, and we hope you are able to have a well-deserved break after another busy year.

Thank you for reading our clinical newsletter during its first year. We hope you received some valuable insights and clinical knowledge that you were able to use in your everyday practice.

Going forward in 2019, Pathology Focus will continue full steam ahead, and we will endeavour to provide you with more clinical articles written by our expert pathologists on topics that are useful and interesting to you. If there are any pathology-related subjects that you would find particularly helpful to your practice, please feel free to email us on newsletter@clinicallabs.com.au

Don't forget that if you would like to switch to the electronic version of the newsletter, which is emailed directly to your inbox, then please feel free to subscribe by faxing back the perforated form, or subscribe online at www.clinicallabs.com.au/subscribe

Inflammatory Skin Disorders:

A practical approach to biopsy diagnosis

Dr Jenny Grew

As the weather warms up and we dispense with a few layers of clothing, it is suspicious lesions and malignancies that feature prominently in presentations of skin disorders to general practitioners. All the more reason to keep in mind that inflammatory skin conditions - the dermatoses - may present at any time.


In this article, some common and not-so-common dermatoses are discussed: how they may present clinically, their typical pathological changes, including the level of skin involved - which in turn informs the best approach to taking a biopsy.

For any biopsy of an inflammatory skin condition, the provision of clinical information is essential and should include:

- Description of the lesions: duration and appearance, age of the lesion and any prior treatment. Consider submitting a clinical photograph to the lab along with the specimen.
- Clinical differential diagnoses.
- Medical, including drug, history.

Let's turn our attention to three of the more commonly encountered dermatoses, each of unknown aetiology.

LICHEN PLANUS



CLINICAL FEATURES

- Typically presents in patients 30-60 years
- Violaceous, flat-topped papules, often polygonal, streaked with white lines.
- Occurs on flexor surfaces of wrists, thighs, genitalia, trunk.
- Oral lesions may occur (in up to 60%)


HISTOLOGY

- Lichenoid pattern:**
- Band-like upper dermal inflammatory infiltrate.
 - Apoptotic keratinocytes in epidermis.
 - Hyperkeratosis and wedge-shaped hypergranulosis.

APPROACH TO BIOPSY

- Incisional biopsy to include superficial dermis
- Include a small amount of normal skin at advancing edge of lesion.
- Punch biopsy in the centre of a lesional area may be suitable.

GRANULOMA ANNULARE



CLINICAL FEATURES

- Typically occurs in patients younger than 30 years.
- More common in women and patients with diabetes.
- Lesions occur on hands, feet, extensor limb surfaces.
- Classical lesion comprises small papules with an enlarging ring configuration.


HISTOLOGY

- Granulomatous: necrobiotic/interstitial.
- Increased dermal mucin.

APPROACH TO BIOPSY

- Incisional biopsy at advancing edge, to include reticular dermis.
- Annular lesions may be unsuitable for punch biopsy.

DISCOID LUPUS ERYTHEMATOSUS



CLINICAL FEATURES

- Peak onset in fourth decade, more commonly occurring in women.
- Wide distribution but classical lesions involve head and neck, including butterfly malar rash.
- Red, scaly, well-demarcated patches with follicular plugging.
- Photo-exacerbation may occur.

HISTOLOGY


- Lichenoid reaction involving epidermis and epithelium of plugged follicles.
- Inflammation involves superficial and deep dermis.
- Direct immunofluorescence may be positive (lupus band test) but not required for diagnosis.

APPROACH TO BIOPSY

- Incisional biopsy to include superficial dermis
- Include a small amount of normal skin at advancing edge of lesion.
- Punch biopsy in the centre of a lesional area may be suitable.

Two Less Common Conditions are Grover Disease and Porokeratosis.

POROKERATOSIS



CLINICAL FEATURES

- Various forms including familial (beginning in childhood), sporadic or associated with immunosuppression.
- May occur on sun-exposed sites (often legs) in middle aged adults as disseminated superficial actinic porokeratosis (DSAP). Lesions may be up to 10mm and resemble solar keratoses.
- Squamous cell carcinoma may develop (except in punctate form) indicating pre-malignant potential of porokeratosis.


HISTOLOGY

- Cornoid lamella is the defining feature: a thin column of parakeratotic cells. Associated with focal loss of the granular layer and with vacuolated and dyskeratotic keratinocytes. Often a lichenoid reaction.

APPROACH TO BIOPSY

- Incisional biopsy at advancing edge, to include reticular dermis.
- Annular lesions may be unsuitable for punch biopsy.

GROVER DISEASE (TRANSIENT/PERSISTENT ACANTHOLYTIC DERMATOSIS)



CLINICAL FEATURES

- Itchy grey-pink papules or papulovesicles, acute eruption.
- Precipitated by sweat, sun exposure, ionising radiation, some drugs.
- Typically occurs on trunk area of middle-aged and older Caucasian men.
- Transient and more persistent chronic forms.

HISTOLOGY

- Tissue reaction pattern of "acantholytic dyskeratosis" with suprabasal clefting.
- Perivascular lymphohistiocytic infiltrate in superficial dermis.
- A few eosinophils may be present.

APPROACH TO BIOPSY

- The changes in Grover disease are superficial, so may be amenable to punch biopsy or shallow incisional biopsy. A solitary papule may be encompassed by a punch sample which ideally would include a surrounding rim of non-lesional skin.

Additional considerations:

Select an appropriate lesion (biopsy at different stages of a lesion according to clinical differential diagnosis):

- an early lesion in itchy or blistering conditions
- an lesion of intermediate duration if vasculitis is suspected
- a late lesion in lupus, psoriasis or fibrosing lesions
- avoid lesions which are ulcerated or have been treated, for example with topical steroids.

Is ancillary testing required?

- **Infection:** microbiology
- **Blistering, vasculitis:** direct immunofluorescence (DIF)
- **Lymphoma:** submit fresh tissue urgently to lab and/or in suitable transport media; discuss with dermatopathologist.

References

Bolgia JL, Schaffer JV, L Cerroni. Dermatology 4th edition. Elsevier, 2018.
 Calonje E, Brenn T, Lazar A and McKee PH. McKee's Pathology of the Skin, 4th Edition. Saunders Elsevier, 2012.
 Photo credits: DermnetNZ, Pathology Outlines, Prof Raimo Suhonen (Discoid Lupus Erythematosus image)

Harvey NT, Chan J and Wood BA. Skin biopsy in the diagnosis of inflammatory skin disease. Australian Family Physician, Vol 46, No. 5, May 2017.
 Patterson JW. Weedon's Skin Pathology, 4th Edition. Churchill Livingstone Elsevier, 2016

Subscribe to our electronic mailing list

Subscribe to the Clinical Labs mailing list and receive our bi-monthly clinical newsletter, important updates, educational resources and more, delivered directly to your inbox. Simply visit clinicallabs.com.au/subscribe and follow the instructions.

Alternatively, complete the form below, tear along the perforated edge and fax it to Clinical Labs Head Office on **(03) 9538 6733**

Title	Given Name
<input type="text"/>	<input type="text"/>

Surname
<input type="text"/>

Email
<input type="text"/>

Practice Name
<input type="text"/>

Practice Address
<input type="text"/>

Practice Suburb	Post Code
<input type="text"/>	<input type="text"/>

Please tick one of the below:

- General Practitioner
 Specialist
 Medical Centre / Practice Manager

Thank you



About the Author

Dr Jenny Grew

MBChB, FRCPA, AFRACMA
Anatomical Pathologist

Email: jenny.grew@clinicallabs.com.au
Phone: 1300 367 674
Location: Subiaco, Western Australia

Expert Pathologist near you



Dr Carl Bulliard

Anatomical Pathology
BSc (Med), MBBS, Grad Dip Med
(Clin Epid), FRCPA

Phone: (02) 8887 9951
State: New South Wales
Email: carl.bulliard@clinicallabs.com.au

Hormone Testing at Clinical Labs

INSULIN-LIKE GROWTH FACTOR-1 (IGF-1)

- IGF-1, also referred to as somatomedin-C (Sm-C), is a hormone similar in molecular structure to insulin that is mainly produced by the liver in blood. IGF-1 mediates most of the effects of Growth Hormone (GH). It plays an important role in childhood growth and continues to have anabolic effects in adults
- IGF-1 is produced throughout life. The highest rates of IGF-1 production occur during the pubertal growth spurt. The lowest levels occur in infancy and old age.
- Levels of IGF-1 can vary in the circulation depending on variation in the levels of GH, insulin levels, genetic make-up, the time of day, age, sex, exercise status, stress levels, nutrition level and body mass index (BMI), disease state, ethnicity, estrogen status and xenobiotic intake.

WHY DO WE MEASURE IGF-1?

- IGF-1: is a clinical severity marker. It correlates with GH levels in the blood so it can be a useful marker in the diagnosis of clinical conditions such as acromegaly and GH deficiency.
- IGF-1 can also be useful in assessing patients' nutritional status, dwarfism, and in monitoring patients on treatment for the above conditions or patients on treatment with recombinant IGF-1.

IGF-1 TEST

- IGF-1 test is a Medicare rebateable
- Can be performed on serum or heparinised plasma samples
- IGF1 test is performed on Siemens Immulite® XPI. Validated Age specific Reference Ranges apply.
- Results are available to clinicians within 2-3 business days.

References

Scarth JP (2006). A review*. Xenobiotica. 36 (2-3): 119-218.
Wade N (17 February 2011). New York Times.
Giustina A, et al., (2014). Nat Rev Endocrinol. 10 (4): 243-8.
Rosenbloom AL (2007). Curr. Opin. Pediatr. 19 (4): 458-64.
Hernandez R (July 2016). Pharm Tech.
Sattler FR (August 2013). Best Pract. Res. Clin. Endocrinol. Metab. 27 (4): 541-55.
Siemens Ref. IMC 18-November 2017



About the Author

A/Prof Mirette Saad

MBBS (Hons), MAACB, FRCPA, PhD
Chemical Pathologist

Email: mirette.saad@clinicallabs.com.au
Phone: (03) 9538 6777
Location: Victoria

ANTI-MÜLLERIAN HORMONE PLUS (AMH PLUS):

As part of the wide range of reproductive pathology services, Australian Clinical Labs has been offering the Anti-Müllerian Hormone (AMH) test using a fully automated Roche Cobas Elecsys® assay which shows excellent analytical performance over other methods including precision, accuracy and functional sensitivity. AMH is an established biomarker produced by antral and pre-antral follicles for assessing ovarian reserve which is considered an important tool in assessing potential fertility.

WHAT'S NEW?

Australian Clinical Labs has introduced the new precise, reliable and robust AMH Plus Immunoassay from Roche Cobas Elecsys®. AMH PLUS enables clinicians to use results when dosing in women undergoing an assisted reproductive technology programme with the human recombinant follicle stimulating hormone (human rFSH), REKOVELLE® (follitropin delta). AMH Age-Specific Reference Ranges (10th-90th percentile) are reported as provided by Roche.

AMH PLUS:

The test is performed on a gel serum tube and clinicians will receive full report of AMH with age specific reference ranges within 2-4 business days. This service is not rebateable by Medicare, and therefore will attract an out of pocket fee.

References

Deeks ED. Elecsys® AMH assay: a review in anti-Müllerian hormone quantification and assessment of ovarian reserve. Mol Diagn Ther 2015; 19: 245-249.
Roche Diagnostics. Elecsys® AMH (anti-Mullerian hormone): Method sheet. 2015. <https://pim-eservices.roche.com>. Last accessed November 2016.
Gassner D, Jung R. First fully automated immunoassay for anti-Müllerian hormone. Clin Chem Lab Med. 2014;52(8):1143-52.
Anderson RA, Anckaert E, Bosch E, et al. Prospective study into the value of the automated Elecsys antimüllerian hormone assay for the assessment of the ovarian growing follicle pool. Fertil Steril. 2015;103(4):1074-80.e4.
Nelson SM, Pastuszek E, Kloss G, et al. Two new automated, compared with two enzyme-linked immunosorbent antimüllerian hormone assays. Fertil Steril. 2015 Oct;104(4):1016-1021.e6.
Hyldgaard J, Bor P, Ingerslev HJ, et al. Comparison of two different methods for measuring anti-müllerian hormone in a clinical series. Reprod Biol Endocrinol. 2015 Sep 22;13(1):107.

To receive our bi-monthly clinical newsletter, updates, educational resources and more, go to clinicallabs.com.au/subscribe and follow the instructions.